

Development of Poly Herbal Mixture from Some Indigenous Plants for Screening of Anti-Microbial Activity.

K.Sudheer Kumar^{1*}, N. Ravindra¹, Hareesh Dara²,

1*. Research Scholar, Pacific Academy of Higher Education and Research University,
Udaipur, Rajasthan, India.

1. MAK College of Pharmacy, # 6 -195/1, Beside JBIT Moinabad Hyderabad - 501504,
Telangana, India.
2. SREE College of Pharmacy, Kothagudem, TS.

Abstract:

Plants contain a diverse group of highly valuable and readily available resource of bioactive metabolites *viz.* Alkaloids, tannins, essential oils and flavonoids. According to World Health Organization has estimated that about 80% of the population in developing countries are unable to afford drugs and rely on traditional medicines especially those that are plan based such as India, Srilanka, Bangladesh, China and Japan. Exploration of the chemical constituents of the plants and pharmacological screening may provide us the basis for developing the lead for development of novel agents. Herbs have provided us some of the very important life savings drugs used in armamentarium of modern medicine, present study helps to know that poly herbal formulations are safe and effective and show synergetic actions. There is a much research is in need for future prospects in drug development and dosage design from natural sources having Antimicrobial activity.

INTRODUCTION

Plants are used as medicine since time immemorial. India is a rich source of medicinal plants. They are widely used in ancient systems of medicine. It is reported that, two or three antibiotics that are launched every year are derived from microorganisms. Antimicrobial activity is the ability of a substance to inhibit or kill bacterial and fungal growth. Different types of antimicrobial and chemotherapeutic agents are being used in the treatment of one form of disease or the other. Herbal medicine is the mainstay of about 75–80% of the world population, mainly in the developing countries, for primary health care because of better cultural acceptability, better compatibility with the human body and lesser side effects. In this present

study an attempt was made to evaluate the Antimicrobial properties of selected indigenous medicinal plants for polyherbal formulation based on the literature studies.

Plant material

Leafs of *Aerva lanata* Plant collected from Urlugonda (village), Suryapet District, and Telangana. Leafs of *Bauhinia variegata* collected from Herbal Garden Chilkur Balaji College of Pharmacy, *Spilanthes acmella* obtained from Herbal garden, Himayath Sagar, Ranga Reddy district , Hyderabad, all the three plants ere authenticated and Botanical identification of the plants was done by Botanical Survey of India, Deccan Regional Centre , Hyderabad, Specimens of *Aerva lanata*, *Bauhinia variegata*, *Spilanthes acmella* (Voucher number:BSI / DRC / 16-17 / Tech/1005).On 22/03/2017.

Drying of plant material

Plant materials were shade dried about for 15-20 days. The shade dried plant material was further coarsely powdered and the powder was passed through the mesh and stored in airtight container for further analysis.



Fig: 1 Powders of *Aerva lanata*, *Bauhinia variegata*, *Spilanthes acmella*

Chemicals and instruments: Compound microscope, glass slides, cover slips, watch glass and other common glasswares and the basic apparatus and instruments, Soxhlate apparatus used for the study. Solvents viz. Methanol, petroleum ether, N-hexane, Ethanol, and reagents viz. phloroglucinol, glycerin, hydrochloric acid, chloral hydrate, Safranin and Fast green. Different strains of microorganism included for study were gram positive bacteria like *Staphylococcus aureus*, *Bacillus subtilius*, gram negative bacteria like *Escherichia coli*, *Klebseilia pneumonia*.

Collection of Bacterial strains

The lyophilized form of different strains of microorganism included for study were gram positive bacteria like *Staphylococcus aureus*, *Bacillus subtilius*, gram negative bacteria like *Escherichia coli*, *Klebseilia pneumoniae*. The bacteria were grown in the nutrient broth at 37°C and maintained on nutrient agar slants at 4°C. All the bacterial strains were procured from

Osmania University, Hyderabad, Telangana. After receiving strains were sub-cultured by using nutrient agar media for evaluation of antimicrobial activity at Microbiology lab at Chilkur Balaji College of Pharmacy.

Phyto-chemical screening

The different extracts were subjected to preliminary Phyto-chemical screening techniques provided in literature


In vitro anti-Microbial activity

The in vitro antibacterial study was performed by measuring the diameter of the zone of inhibition on the inoculum agar plate. The zone of inhibition can be defined as the clear region around the susceptible disc or well with an antimicrobial agent on the agar surface. It is designated to test the ability of the antimicrobial agent to inhibit the growth of microorganisms [150]. The larger the zone of inhibition is, the better the inhibition of the particular antimicrobial agent.

Fig 02 : Inhibition zones of A) *Aerva lanata* , B) *Bauhinia variegata* , C) *Acmella uliginosa*

Preparation of Nutrient agar plates

28 g of nutrient agar was dissolved in 1000 ml of distilled water and boiled to dissolve the agar completely and then sterilized by autoclaving at 15 lbs pressure at 121°C for 15 minutes (pH



7.4±0.2). 20-25 ml of cooled agar medium was poured onto the sterile prepared 15 × 100 mm sterile Petri dishes. (The agar plates were allowed to cool to room temperature and stored in a refrigerator (2°C - 8°C) until used) Evaluation of anti-Microbial activity by Agar Well Diffusion

Evaluation of Antibacterial activity:

Leaf extracts of *Aerva Lanata*, *Bauhinia variegata*, *Acmella uliginosa* at a concentration of 500µg/ml, 750µg/ml, 1000µg/ml were tested against the gram positive bacteria like *Staphylococcus aureus*, *Bacillus subtilis*, gram negative bacteria like *Escherichia coli*,

Klebsiella pneumoniae, by Well Diffusion Method, the same method employed for evaluation of antifungal activity with some modifications in temperatures and standard drug according to literature survey and standard methods employed previously.

Antibacterial activity of the plants extract was tested using Well diffusion method.³¹ The prepared culture plates were inoculated with different selected strains of bacteria using streak plate method. Wells were made on the agar surface with 6mm cork borer.

The dried extracts were dissolved in 95% of DMSO for preparation of different concentration ranges of extracts. The extracts were poured into the well using sterile syringe. The plates were incubated at $37^{\circ}\text{C} \pm 2^{\circ}\text{C}$ for 24 hours for bacterial activity. The plates were observed for the zone clearance around the wells. The extracts of the dried scale leaves of three plants were used for the study. The extracts were dissolved in DMSO to form dilution such as $500\mu\text{g/ml}$, $750\mu\text{g/ml}$ and $1000\mu\text{g/ml}$. Each concentration of the extract was tested against different bacterial pathogens. Ciprofloxacin, at a concentration of $5\mu\text{g/ml}$ and $10\mu\text{g/ml}$ was used as standard antibacterial drug. The zone of inhibition was calculated by measuring the diameter of the inhibition zone around the well (in mm) including the well diameter. The readings were taken in three different fixed directions in all three replicates³² and the average values were tabulated. Table No : 13

Triplicates were performed and the experiments were repeated thrice and the average values were recorded. The results were compared with the zone of inhibition produced by standard anti microbial drug.

From the above results came to know that the selected plants shown significant antibacterial activity in methanolic extract, in individual, then we made an attempt for evaluation of synergistic effect of selected medicinal plants leafy extracts of *Aerva Lanata*, *Bauhinia variegata*, *Acmella uliginosa*, and determined the study of antimicrobial activities against same species of bacterial strains with same procedure employed above i, ie Agar Well Diffusion and results were reported at Table No: Table No: 13,14,15

RESULTS**Table No: 1****1. Morphological Evaluation**

Parameter	<i>Aerva lanata</i>	<i>Bauhinia Variegata</i>	<i>Spilanthes acmella</i>
Color	Pale green	Greenish	Dark green
Odor	Characteristic	Weak	Strong
Taste	Bitter to acrid	Slightly bitter	Burning, pungency, numbness
Shape	Ovate	Cordate, nerved	ovate,narrowed at base
Texture	Smooth	Rough	Smooth
Size	0.5 to 1.5 in wide (13 to 38 mm) long	13 to 15 cm length 2-14 cm wide	2.5 to 5 cm long 1.5 to 2 cm wide

Table No:- 2 phytochemical evaluations of *Aerva lanata* leaves

Sl. No	Name of extract	Alkaloids	Sapoinins	Proteins	Glycosides	Flavonoids	Carbohydrates	Phyosterols	Tannins
1	N-hexane	+	+	-	-	+	+	+	-
2	Petroleum ether	+	+	-	-	+	+	-	-
3	Ethanol	+	+	+	-	+	+	+	
4	Methanol	+	-	-	-	+	+	-	+

Table No:- 3 Phytochemical evaluation of *Bauhinia variegata*

Sl. No.	Name of extract	Alkaloids	Sapoinins	Pro teins	Glycosides	Flavonoids	Carbohydrates	Phytosterols	Tannins
1	N-hexane	-	+	+	-	+	+	+	-
2	Petroleum ether	-	+	-	-	+	+	+	+
3	Ethanol	-	+	+	-	+	+	+	
4	Methanol	-	+	-	+	+	+	+	+

Table No:- 4 phytochemical evaluation of *Spilanthes acmella*

Sl. No.	Name of extract	Alkaloids	Sapoinins	Pro teins	Glycosides	Flavonoids	Carbohydrates	Phytosterols	Tannins
1	N-hexane	+	+	+	-	+	+	-	+
2	Petroleum ether	+	+	-	-	+	+	-	+
3	Ethanol	+	+	+	-	+	+	-	+
4	Methanol	+	-	-	-	+	+	-	+

***Where, + indicates the presence of secondary metabolites, - indicates the absence of secondary metabolite

Anti microbial evaluation

Table No:- 5 Results for Anti-Bacterial activity of *Aerva lanata* leaves

Solvent extracts 1000µg/ml	Ciprofloxacin		Zone of Inhibition			
	5µg/ml	10µg/ml	Staphylococcus aureus,	Bacillus subtilius	Escherichia coli,	Klebseilia pneumoniae
N-hexane	7.5 mm	9 mm	5 mm	6 mm	7 mm	5 mm
Petroleum ether	7 mm	9 mm	6 mm	7 mm	6 mm	5 mm
Ethanol	7 mm	9 mm	7 mm	6 mm	7 mm	8 mm
Methanol	7.5 mm	9.5 mm	8 mm	6.5 mm	8 mm	8 mm

Table No:- 6 Results for Anti-Bacterial activity *Bauhinia variegata*

Solvent extracts 1000µg/ml	Ciprofloxacin		Zone of Inhibition			
	5µg/ml	10µg/ml	Staphylococcus aureus,	Bacillus subtilius	Escherichia coli,	Klebseilia pneumoniae
N-hexane	7.5 mm	9 mm	3 mm	4mm	5 mm	4 mm
Petroleum ether	7 mm	6 mm	4 mm	3 mm	4 mm	3 mm
Ethanol	7 mm	7 mm	5 mm	4 mm	5 mm	5 mm
Methanol	7.5 mm	9.5 mm	9 mm	8mm	7 mm	9 mm

Table No:- 7 Results for Anti-Bacterial activity *Spilanthes acmella*

Solvent extracts 1000µg/ml	Ciprofloxacin		Zone of Inhibition			
	5µg/ml	10µg/ml	Staphylococcus aureus,	Bacillus subtilius	Escherichia coli,	Klebseilia pneumoniae
N-hexane	7.5 mm	9 mm	4 mm	4mm	6 mm	4 mm
Petroleum ether	7 mm	6 mm	5 mm	4 mm	5 mm	6 mm
Ethanol	7 mm	7 mm	6 mm	5 mm	6 mm	7 mm
Methanol	7.5 mm	9.5 mm	10 mm	9 mm	9 mm	8 mm

DISCUSSION

Literatures revealed that the selected three herbs *Aerva lanata*, *Bauhinia variegata*, *Spilanthes acmella* have antibacterial activity. Extraction and the phytochemical screening was done using Methanol, petroleum ether, N-hexane, and Ethanol as solvents. Phytochemical screening confirmed the presence of various phytoconstituent like carbohydrate, glycosides, flavonoids, Alkaloids, Volatile oils and tannins. In the present study, polyherbal ointments were prepared by fusion method using emulsifying ointment as the base. Antimicrobial study shows that the prepared ointments has better and equal activity against gram positive bacteria like *Staphylococcus aureus*, *Bacillus subtilius*, gram negative bacteria like *Escherichia coli*, *Klebseilia pneumonia*, compared to standard 5% Betadine

CONCLUSION

. From this study concluding that poly herbal formulations are safe and effective and show synergetic actions. There is a much research is in need for future prospects in drug development and dosage design from natural sources.

REFERENCES

1. Amutha K, Antimicrobial activity of *Aerva lanata*, *Boerhavia diffusa*, *Cassia tora*, *Pongamia pinnata* and influence on the virulence factors of multi-drug resistant *Escherichia coli* and *Pseudomonas aeruginosa*, U publish.info, 2010:1
2. Chowdary D , Sayeed A et al, Antimicrobial activity and cytotoxicity of *Aerva lanata*, *Fitoterapia*, 2002;73(1):92-4.
3. Muthukumaran P, Shanmuganathan P, et al, Antioxidative and Antimicrobial Study of *Aerva Lanata*, *Asian Journal of Biochemical and Pharmaceutical Research*, 2011; 2 (1):265-67
4. Gurumurthy H et al, *In vitro* antibacterial activity of *Aerva lanata* (L.) Juss, *Environment and Ecology*, 2009; 27(1): 32-35
5. John De Britto A, et al, Antibacterial activity of a few medicinal plants against *Xanthomonas campestris* and *Aeromona hydrophila* , *Journal of Biopesticides*, 2011; 4 (1): 57 - 60
6. Surendra KP et al, Review on Herbal therapy for Antiurolithiatic activity, *Journal of Pharmacology and Toxicology*, 2011;6: 321-32.
7. Bhatia L, Bishnoi H, Chauhan P, Kinja K, Shailesh S. *In-vitro* comparative antioxidant activity of ethanolic extracts of *glycosmis pentaphylla* and *bauhinia variegata*. *Recent Res Sci Technol*. 2011;3:01–3. (XX1)
8. Umachigi SP, Jayaveera K, Ashok Kumar CK. Evaluation of wound healing potential of poly herbal formulation. *Pharmacol Online*. 2009;3:505–12.(XX2)
9. Nevin K.G., Effect of *Aerva lanata* against hepatotoxicity of carbon tetrachloride in rats, *Environmental Toxicology and Pharmacology*, 20(3); 2005, 471-477.
10. Pathiratna L S S, Joseph K.D.S.M. and Perera M.K.P., The effect of some cultural practices on the growth and yield of the medicinal plant *Aerva lanata* (L.) Juss. *Ex schult. (polpala)*, *Cey. J. Sci. (Bio. Sci.)* 32;2004, 67-74.
11. Akram A, Abdellatif N, Mohamed F N. Urolithiasis in Tunisian children: A study of 100 cases. *Soudi J Of Kidney Disease and Transplantatio*, 2009; 2(6): 1096-1100.
12. Nevin KG, Vijayammal PL, Effect of *Aerva lanata* on solid tumor induced by DLA cells in mice. *Fitoterapia*, 2003; 74:578.
13. Vetrichelvan T, Jagadeesan M, Palaniappan MS, Murli NP. and Sasikumar K. Diuretic and anti-inflammatory activities of *Aerva lanata* in rats. *Indian Journal of Pharmaceutical Sciences*, 2000;62: 300-302.